

Neurodynamics of vertebrogenic somatosensory activation and Autonomic Reflexes - a review:

Part 3 A central connection

Peter Rome and John Waterhouse

Abstract: A review is presented of the central nervous system association with noxious sensory activation of vertebral articulations from disturbed segmental mechanics. Examples of neurological studies reflecting and supporting this physiological function are presented.

Indexing terms: Vertebrogenic; somatosensory; autonomic reflexes; subluxation; chiropractic

Introduction

An early example of the association between vertebral dysfunction and the brain was reported in 1958 by the European medical doctor Sollman. He noted functional changes in the diencephalon through manipulation of the upper cervical spine influencing sympathetic afferents and cardiac physiology. One third of his 150 patients showed positive vegetative changes. (1, 2, 3, 4, 5)

Carrick established an innovative paradigm of functional clinical neurology in a specialised and systematic extension of chiropractic. This incorporated a model of analysis, diagnosis and management involving the brain and the autonomic nervous system (ANS). This model of neurology seeks to perceptively interpret a range of traditional as well as abstruse and subtle clinical signs, to reveal an elaborate diagnostic neurological disturbance which may influence therapeutic direction and management. (6, 7)

Instances of broad vertebrogenic influence on nociception were noted by Sparks et al, who monitored this association by the reduction of pain through thoracic spinal manipulation and demonstrated functional MRI (fMRI) changes in cerebral circulation. (8, 9)

An analgesic effect was noted when specific regions of the brain were highlighted in another fMRI study following somatic stimulation through joint mobilisation of the hind limb of rats. Malisza et al noted decreased areas of activation bilaterally in the anterior cingulate and frontal cortex, as well as in the contralateral sensory motor cortex. Further somatosensory-autonomic association through the sympathetic nervous system was explored via fMRI studies by Henderson and Macefield in a 2013 report. (10, 11)

... Mechanical disturbance of vertebral articulation can affect proprioception through the ANS as dyskinesthesia. This proprioceptive disturbance may be monitored in clinical testing. Evidence also indicates that such ANS disturbances may be moderated by addressing the vertebral dysfunction of the somatic-mechanical input, through manual corrective spinal manipulation.'



Also using fMRI imaging, a randomised placebo controlled study by Cerritelli et al provided an insight into brain activity through osteopathic manual therapy. (12, 13)

Also monitoring somatosensory evoked potentials and sensorimotor integration, Haavik-Taylor and Murphy recorded short interval intracortical facilitation, as well as inhibition, and cortical silent periods, in response to manipulation of cervical joint dysfunction (vertebral subluxations) in humans. A later study suggested that the procedure can lead to transient cortical plastic changes. (14, 15, 16)

A 2005 Russian study of 134 children who had previously experienced cervical trauma, correlated a spinoneural connection between segmental spinal excitability of pain with polysynaptic brain excitability as monitored by spino-bulbo-spinal, blink, and H-reflexes. (17)

An Interdisciplinary Spinal Research group formed in 2016 through the *Department of Chiropractic Medicine at Balgrist University Hospital* in Switzerland was comprised of '*multiple medical disciplines (chiropractic, medicine, physiotherapy, molecular biology, movement sciences and neuroscience)*'. They investigated several aspects of lower back pain including the wider ramifications from biomechanical somatosensory influences reflected in cortical representation and associated brain circulatory changes. (18, 19, 20, 21)

An extensive examination of the close association of the somatovisceral and viscerosomatic association with spinal manipulation and neural integration was conducted at a symposium organised by the American Academy of Osteopathy in 1989. In considering the evidence presented, the editors stated that '*the information being collected in laboratories around the world is relevant to the clinical experiences of the osteopathic profession*'. (22)

Somatosensory Centralisation *Sensory Evoked Potential and Sensorimotor*

Somatosensory centralisation into the CNS is an aspect of the integrated reactivity of the autonomic nervous system in the presence of persistent pain. This results in heightened sensitivity to pain, and even to light touch. Manual therapists seek to remove that initiating stimulus of the pain, in order to normalise the noxious stimulation of the ANS. While they can be generated by mechanical stimulation, electrical activations are often used in studies which appear to induce a more robust response. (23, 24, 25, 26, 27, 28)

Central sensitization (CS) is essentially an enhancement within the CNS - especially of activated reflex circuits attributed to noxious stimulation. It includes recruitment of subthreshold synapses and neural plasticity resulting in an augmented stimulation of somatosensory input. Latremoliere and Woold state that CS is '*responsible for many of the temporal, spatial, and threshold changes in pain sensibility in acute and chronic clinical pain settings, and exemplifies the fundamental contribution of the central nervous system to the generation of pain hypersensitivity*.' (29, 30, 31, 32)

Prominent researchers in this field - Haavik, Murphy, Holt, and others, have utilised the somato-autonomic-visceral CNS pathways to monitor Somatosensory Evoked Potential (SEP) and sensorimotor changes, and their integration with spinal manipulation. (33, 34, 35, 36, 37, 38, 39, 40, 41, 42)

In further integration, Cersosimo and Benarroch note that the central autonomic networks '*are reciprocally interconnected (and) receive converging visceral and somatosensory information (that) generate stimulus specific patterns of autonomic, endocrine, and motor responses*.' (43)

Zusman noted a significant role for manual therapy, in that it may impact upon sudomotor function, heart rate, blood pressure and plasma hormones through influencing the ANS. He found that pain inhibition was through the manipulative effect of activating descending brain stem inhibitory pathways. (44)

As SEPs and sensorimotor reflexes involve brain connectivity, they can be monitored by electromagnetic evaluation which can be reflected in EEG changes. It was noted that abnormal SEPs can arise from dysfunction of a range of innervated structures such as spinal articulations, nerve roots and the trigeminal nerve. They can also involve afferent neural tracts through the periaqueductal gray region (PAG) where one of its roles is nociceptive inhibition. (45, 46, 47, 48, 49)

As well as relieving a somatic irritant, a 1995 study by Wright indicated that cervical mobilisation may activate descending inhibitory nociceptive pathways from the dorsal periaqueductal gray area of the midbrain (dPAG). These findings were supported by Sterling and colleagues in 2001. They noted excitation of the sympathetic nervous system, as well as a contrary response between the deep and superficial neck flexors. This concept has subsequently been noted by Haavik and colleagues and also by Falco and colleagues, through cervical facet anaesthetic blocks. (14, 15, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61)

In 1984 after labelling both the splanchnic as well as the intercostal nerves, Pierau and colleagues were able to demonstrate somatovisceral neural integration in a cat. They demonstrated splanchnic and intercostal nerve convergence in the dorsal root ganglia, and through the ventral and lateral horns of the lower dorsal spinal cord. (62)

Kimura and Sato state that *'Somatic sensory information from the skin, muscles and joints is transmitted to the central nervous system (CNS) by somatic afferent nerves and induces sensation, emotion and reflex responses of motor and autonomic functions.'* They also proposed that somato-autonomic reflex responses tend to explain the efficacy of spinal manipulation, physiotherapy, acupuncture and moxibustion. (36, 42, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74)

In a physiotherapy study, Bialosky et al suggest that manual manipulative intervention *'initiates a cascade of neurological responses ... which are then responsible for the clinical outcomes.'* (75)

Cerebellum

Functional cerebellar changes associated with manipulative care have also been noted. In 2013, Daliqadu and colleagues interpreted readings from transcranial magnetic stimulation of the cerebellum following neck manipulation. They monitored motor sequence learning interventions to demonstrate a decrease in cerebellar inhibition and improved task performance. They enlisted 10 subclinical neck pain subjects and compared the findings with 10 healthy controls in the study. (76)

Haavik and Murphy demonstrated that even subclinical neck pain can affect joint position sense of the elbow – then further, that position accuracy improved following adjustments of the cervical subluxations. (33, 35, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87)

Thalamus

An indication of the influence of spinal manipulation on the rat thalamus was reported by Reed and colleagues in 2014 and again in 2017. This collaborative medical-chiropractic research indicated that manipulation of the lumbar spine may suppress *submedius* and *lateral neuronal activity* in the thalamus, where it may be a factor in pain modulation before nociceptive input reaches the cortex. Similar research was undertaken by others, and found to affect thalamic regions following stimulation from either acupuncture or morphine.

Other central connections of somatovisceral reflexes may involve neural plasticity with the thalamus. For instance, in 1985 a study by Vaganian et al demonstrated somatovisceral integration through somatic afferent stimulation (in cats). They found it to be inhibitory and potentially involved the thalamus. (3, 88, 89, 90)

Nociception

Nociceptive input to the ANS constitutes interoception as a physiological condition of the entire body, not just the viscera. Amongst other etiological factors to consider with this are the duration, severity, and anatomical location of the noxious input. In addition, research as to the chronicity of subtle (mild) nociceptive neural irritation upon target structures over extensive periods of time, may provide insight into the development of clinical signs of organ dysfunction which in turn may eventually lead to pathological tissue changes. (91, 92, 93)

Rather extensive autonomic changes have been noted resulting from musculoskeletal pain. These include; autonomic dysregulation, altered pain processing, modulation involving the adrenergic pathway, and catecholamine responses. A physiological response to somatic pain has been shown to affect heart rate, blood pressure, and heart rate variability. (94, 95, 96, 97)

Pain of mechanical spinal origin or degrees of it may be the most common source of aberrant sensory input in somato-autonomic reflexes. As such, mechanical back pain must be a frequent stimulator of somatosensory input to the ANS, affecting other activated ANS reflex functions of innervated structures.

A further ANS irritant could be the chemical and thermal nociceptive input of inflammation involving disturbed articulations, commonly presented for manipulative therapy. As a protopathic mechanism, it becomes an important and convenient diagnostic aid. As pain activates somatosensory and somatovisceral reflex pathways, its removal would represent a *raison d'être* for the manipulative sciences. Budgell states that *'Pain in general, and perhaps spinal pain in particular, is capable of eliciting changes in visceral function which can be distressing and even dangerous.'* (98, 99)

Relief from spinal pain including cervicgia, headaches, and radicular pain extending from the spine, are two of the more common conditions addressed by manipulation. The neurology of this pain however is heterogeneous. It is a further example of somato-autonomic interface as to the potential influence of spinal manipulation. Benarroch states that *'the interaction between the nociceptive and the autonomic systems are complex, and involve a variety of central antinociceptive, autonomic, emotional, and behavioural control mechanisms...and could provide the basis for innovative pharmacological, physical, and behavioural therapy approaches.'* (100, 101, 102, 103, 104)

In light of recent concerns over opioid addiction, it is possible that greater attention should be afforded to the physical, and behavioural model of pain management. This could be in the form of conservative initial triage for analgesic care and its broader somatosensory ramifications. (105, 106)

The analgesic effects of manual therapy were also reviewed by Vigotsky and Bruhns in 2015. They noted that different forms of manual therapy resulted in different analgesic effects and different mechanisms, but all elicit a neurophysiological response. (107)

If somatic nociceptive activation from disturbance of vertebral articulations can activate ANS reflexes, they may then have the potential to adversely affect visceral function through noxious somatovisceral reflexes. It would then seem feasible to expect that removal of the noxious stimulus to ameliorate or moderate the activated pathophysiologic somatovisceral reflexes would lead to a normalisation of those same reflexes. Burton and colleagues state that *'The sympathetic nervous system is inherently involved in a host of physiological responses evoked by noxious stimulation. Experimental animal and human models demonstrate a diverse array of heterogeneous reactions to nociception.'* (108, 109)

Proprioception - sensorimotor

Proprioception (kinesthesia) provides one of the body's most extensive sensory inputs for position sense and movement. It provides both conscious and subconscious position awareness, and is protopathic in nature. Mechanical disturbance of vertebral articulation can affect proprioception through the ANS - dyskinesthesia. This proprioceptive disturbance may be monitored in clinical testing. Evidence also indicates that such ANS disturbances may be moderated by addressing the vertebral dysfunction of the somatic-mechanical input, through manual corrective spinal manipulation.

Proprioception afferent input is supplied from Group I and II sensory neurons. Posturally, there is critical input of position from these mechanoreceptors in paravertebral muscles and articulations. Optimal proprioception is essential for equilibrium. It is a major consideration in preventing falls, particularly in older patients. Consequently aberrant proprioceptive feedback from disturbed joints may lead to balance and co-ordination problems. (69, 77, 110, 111, 112, 113, 114, 115, 116)

In noting that proprioception is also derived from muscle and other soft tissue structures, as well as vision and vestibular input. Swinkels suggested that proprioception could be disturbed by degenerative spinal articulations, due to the resultant lack of physiological proprioceptive input from the normal articular structures. (117, 118)

Apart from nociception, a noxious somatosensory barrage may also arise from other disturbed mechanoreceptors from involved structures including - *Pacinian corpuscles* (pressure receptors), *Meissner corpuscles* (movement of pressure), *Merkel cells* (light touch), *Golgi tendon organs* (proprioception) and *muscle spindles* (stretch); all combining to register an awareness of proprioception. Chronic stimulation of these structures has been demonstrated to affect ANS activity. Zimny stated that articular tissue comprises *Ruffini-like receptors* (pressure), Golgi tendon organs, and *Pacinian-like corpuscles* (pressure) are found in the joint capsules, ligaments and cartilage, in addition to free nerve endings. These disturbances can be reflected in aberrant proprioception activities. (40, 77, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128)


Sudomotor

Somatosympathetic reflexes may also be monitored through superficial temperature and sweat gland activity, in regions where it correlates with mechanically disturbed segmental spinal influence. (129, 130, 131, 132, 133, 134)

Significantly, and in confirmation of the potential for vertebrogenic conditions, Sato, Sato, and Schmidt revealed that '*the existence of spinal centres for somato-autonomic reflexes had been denied until quite recently*'. (135) General adoption of the possible principles contained in this statement seemed confined primarily to chiropractic and osteopathy.

Conclusion

In essence, the reduction of noxious somatic afference originating from these neural beds in disturbed vertebral facets (subluxation), would seem to have the potential to moderate ANS associated pathophysiological reflex influence which may be linked to visceral dysfunction.



Peter Rome

DC (ret), FICC

cadaps@bigpond.net.au

John D Waterhouse

DC, FACC

Private practice, Melbourne

Cite: Rome P, Waterhouse JD. Neurodynamics of vertebrogenic somatosensory activation and Autonomic Reflexes - a review: Part 3 A central connection. *Asia-Pacific Chiropr J.* 2021;1.4. URL apcj.net/papers-issue-2-4/
#RomeWaterhousePart3CentralConnection

References

1. Sollmann A. Funktionsänderungen des Zwischenhirns durch manipulative Wirbelsäulentherapie. [Functional changes in the diencephalon brought about by manipulative vertebral therapy.] *Hippocrates* 1958;29(7):202-204. (German) (Pubmed extract)
2. Sollman AH. [Disturbed spinal kinematics.] *Zbl Chir* 1960;85:1154-60. (German)
3. Cauwenbergs P. Vertebral subluxation and the autonomic nervous system. In: Gatterman MT Ed. *Foundations of chiropractic subluxation*. 2nd ed. St Louis, Elsevier; 2005;305-336
4. Patterson MM, Howell JN. Editors. *The central connection: somatovisceral/viscerosomatic interaction*. Proceedings of the 1989; Am Osteop Acad Symposium; Athens, Ohio. University Classics. 1992.
5. Margach RW. Chiropractic functional neurology: an introduction. *Integr Med (Encinitas)*. 2017;16(2):44-45.
6. Carrick FR. Changes in brain function after manipulation of the cervical spine. *J Manipulative Physiol Ther* 1997;20(8):529-545. <https://carrickinstitute.com/>
7. Daubeny N, Carrick FR, Melillo RJ, Leisman G. Effects of contralateral extremity manipulation on brain function. *Int J Disabil Hum Dev*. 2010;9(4):269-273.
8. Sparks C, Cleland. Elliot JM, Liu W-C. Using functional magnetic resonance imaging to determine if cerebral hemodynamic responses to pain challenge following thoracic spine thrust manipulation in healthy individuals. *J Orthop Sports Phys Ther*. 2013;43(5):340-8.
9. Sparks C, Liu WC, Cleland JA, et al. Functional magnetic resonance imaging of cerebral hemodynamic responses to pain following thoracic thrust manipulation in individuals with neck pain: a randomised trial. *J Manipulative Physiol Ther*. 2017;40(9):625-634.
10. Malisza KL, Gregorash L, Turner A, et al. Functional MRI involving painful stimulation of the ankle and the effect of physiotherapy joint mobilisation. *Magn Reson Imaging*. 2003;21(5):489-496.
11. Henderson LA, Macefield VG. Functional imaging of the human brainstem during somatosensory input and autonomic output. *Frontiers Human Neurosci*. 2013;7:1-8
12. Cerritelli F, Chiacchiaretta P, Gumbi F, et al. Effects of manual approaches with osteopathic modality on brain correlates of interoception: an fMRI study. *J Vert Subluxation Res*. 2020; May 28:79-92
13. Gyer G, Inklebarger MJ, Tedla J. Spinal manipulation therapy: is it all about the brain? A current review of the neurophysiological effects of manipulation. *J Integr Med*. 2019;17(5):328-37. doi: 10.1016/j.joim.2019.05.004.
14. Haavik-Taylor H, Murphy B. Cervical spine manipulation alters sensorimotor integration: a somatosensory evoked potential study. *Clin Neurophysiol*. 2007;118(2):391-402
15. Haavik-Taylor H, Murphy B. Altered sensorimotor integration with cervical spine manipulation. *J Manipulative Physiol Ther*. 2008;31(2):115-126
16. Robinault L, Holobar A, Crémoux S, et al. The effects of spinal manipulation on motor unit behaviour. *Brain Res*. 2021;11(1):105. doi: 10.3390/brainsci11010105. <https://pubmed.ncbi.nlm.nih.gov/33466707/>
17. Liev AA, Scorobogach MI, Kulkovsky BT. Suprasegmental and segmental activity of the cerebrum and spinal brain in myofascial pain syndrome of children with consequences of patrimonial trauma to the cervical spine. *Manual Therapy. [Мануальная Терапия]* 2005. http://www.mtj.ru/mtj_2005_english_summary_1.pdf. (Abstract)
18. Integrative Spinal Research [ISR]. Dept of Chiropr Med, Balgrist University Hospital, Switzerland. <http://painresearch.ch/index.php>
19. Hotz-Boendermaker S, Marcar VL, Meier ML, Boendermaker B, Humphreys BK. Reorganization in secondary somatosensory cortex in chronic low back pain patients. *Spine* 2016; 41(11):E667-73.
20. Vrana A, Meier ML, Hotz-Boendermaker S, Humphreys BK, Scholkmann F. Different mechanosensory stimulations of the lower back elicit specific changes in hemodynamics and oxygenation in cortical sensorimotor areas-A fNIRS study. *Brain Behavior* 2016;6(12):e00575
21. Boendermaker B, Meier ML, Luechinger R, Humphreys BK, Hotz-Boendermaker S. The cortical and cerebellar representation of the lumbar spine. *Human Brain Mapping*. 2014;35(8):3962-3971.
22. Patterson MM, Howell JN. (4). p XI

23. Hobbs SF, Chandler MJ, Bolser DC, Foreman RD. Segmental organisation of visceral and somatic input onto C3-T6 spinothalamic tract cells of the monkey. *J Neurophysiol* 1992;68(5):1575-1588.
24. Patterson MM, Steinmetz JE. Long-lasting alterations of spinal reflexes: A potential basis for somatic dysfunction. *Manual Medicine*. 2: 38-42.
25. McAllister MJ. What is central sensitization? Institute for Chronic Pain. <http://www.instituteforchronicpain.org/understanding-chronic-pain/what-is-chronic-pain/central-sensitization> May 29, 2017
26. Foreman RD. The functional organisation of visceral and somatic input to the spinothalamic system. (4 – p 178-207)
27. Johnston WL. Osteopathic clinical aspects of somatovisceral interaction. (4 – p30-46)
28. Legatt AD, Benbadis SR. General principles of somatosensory evoked potentials. *Medscape. Drugs and Diseases*. Dec, 2014. <https://emedicine.medscape.com/article/1139906-overview>
29. Latremoliere A, Woolf CJ. Central sensitization: a generator of pain hypersensitivity by central neural plasticity. *J Pain*. 2009;10(9):895-926
30. Woolf CJ. Central sensitization: implications for the diagnosis and treatment of pain. *Pain*. 2011;152(3 Suppl):S20S15. Doi: 10.1016/j.pain.2010.09.030
31. Haavik-Taylor H, Murphy B. Cervical spine manipulation alters sensorimotor integration: a somatosensory evoked potential study. *Clin Neurophysiol*. 2007;118(2):391-402.
32. Zaferea J, Deschenes BK. The role of spinal manipulation in modifying central sensitization. *J Applied Biobehav Research*. 2015;20(2): DOI: 10.1111/jabr.12033
33. Baarbé JK, Holmes MW, Murphy HE, Haavik H, Murphy BA. Influence of subclinical neck pain on the ability to perform a mental rotation task: a 4-week longitudinal study with a healthy control group comparison. *J Manipulative Physiol Ther*. 2016;39(1):23-30.subclinical
34. Dishman JD, Ball KA, Burke J. Central motor excitability changes after spinal manipulation: a transcranial magnetic stimulation study. *J Manipulative Physiol Ther*. 2002;25(1):1-9.
35. Haavik H, Murphy B. The role of spinal manipulation in addressing disordered sensorimotor integration and altered motor control. *Electrophys Kinesiol*. 2012;22(5):768-776 used later also
36. Haavik H, Niazi IK, Holt K, Murphy B. Effects of 12-weeks of chiropractic care on central integration of dual somatosensory input in chronic pain patients: a preliminary study. *J Manipulative Physiol Ther*. 2017;40(3):127-138.
37. Haavik-Taylor H, Holt K, Murphy B. Exploring the neuromodulatory effects of the vertebral subluxation and chiropractic care *Chiropr J Aust*. 2010;40(1):37-44
38. Haavik-Taylor H, Murphy B. Altered sensorimotor integration with cervical spine manipulation. *J Manipulative Physiol Ther*. 2008;31(2):115-126
39. Haavik-Taylor H, Murphy B. Cervical spine manipulation alters sensorimotor integration: a somatosensory evoked potential study. *Clin Neurophysiol*. 2007;118(2):391-402 CORTICAL
40. Holt KR, Haavik H, Lee AC, Murphy D, Elley CR. Effectiveness of chiropractic care to improve sensorimotor function association with falls risk in older people: a randomised controlled study. *J Manipulative Physiol Ther*. 2016;39(4):267-278.
41. Lelic D, Niazi IK, Holt K, et al. Manipulation of dysfunctional spinal joints affects sensorimotor ion in the prefrontal cortex: a brain source localization study. *Neural Plast*. 2016;3704964.doi: 10.1155/2016/3704964.
42. Taylor HH, Murphy B. Altered central integration of dual somatosensory input after cervical spine manipulation. *J Manipulative Physiol Ther*. 2010;33(3):178-188.
43. Cersosimo MG, Benarroch EE. Central control of autonomic function and involvement in neurodegenerative disorders. *Handb Clin Neurol*. 2013;117:45-57. doi: 10.1016/B978-0-444-53491-0.00005-5.
44. Zusman M. Mechanism of mobilization. *Physical Ther* 2011;16(4):233-236. Behbehani MM. Functional characteristics of the midbrain periaqueductal gray. *Prog Neurobiol*. 1995;46(6):575-605.
45. Legatt AD, Benbadis SR. General principles of somatosensory evoked potentials. *Medscape*. <http://emedicine.medscape.com/article/1139906-overview>. 2014
46. Anguera JA, Seider RD, Gehring WJ. Changes in performance monitoring during sensorimotor adaption. *J Neurophysiol*. 2009;102(3):1868-1879.
47. Sterling M, Jull G, Wright A. Cervical mobilisation: concurrent effects on pain, sympathetic nervous system activity and motor activity. *Man Ther*. 2001;6(2):72-81.
48. Barwell R, Long A, Byers A, Schisler C. A four case study: the effect of chiropractic adjustment on the brain wave pattern as measured by QEEG. Summarising an additional 100 (approximately) cases over a three year period. *The Chiropr J*. <http://www.torquerelease.com.au/Effect-of-Chiropractic-adjustment-on-brain-wave-pattern.pdf>. 2008
49. Lelic D, Niazi IK, Holt K, et al. Manipulation of dysfunctional spinal joints affects sensorimotor integration in the pre-frontal cortex: as brain source localisation study. *Neural Plasticity*. 2016;3704964. doi: 10.1155/2016/3704964.
50. Budgell B, Sato A. Modulations of functions by somatic nociceptive inputs. *Prog Brain Res*. 1996; 113:535-539.

51. Burton AR, Fazalbhoy A, Macefield VG. Sympathetic responses to noxious stimulation of muscle and skin. *Front Neurol.* 2016;30(7):109. doi: 10.3389/fneur.2016.00109.
52. Cleland JA, Childs JD, McRae M, Palmar JA, Stowell T. Immediate effects of thoracic manipulation in patients with neck pain: a randomized clinical trial. *Man Ther.* 2005;10(2):127-135.
53. Coronado RA, Gay CW, Bialosky JE, et al Changes in pain sensitivity following spinal manipulation: a systematic review and meta-analysis. *J Electromy Kinesiol.* 2012;22(5):752-767.
54. Falco F, Erhart S, Wargo BW, et al. Systematic review of diagnostic utility and therapeutic effectiveness of cervical facet joint interventions. *Pain Physic* 2009;12(2):323-344.
55. Fazalbhoy A, Birznies I, Macefield VG. Individual differences in the cardiovascular responses to tonic muscle pain: parallel increases or decreases in muscle sympathetic nerve activity, blood pressure and heart rate. *Exp Physiol.* 2012;97(10):1084-1092.
56. Jinkins JR, Whittemore AR, Bradley WG. The anatomical basis of vertebral pain and autonomic syndrome associated with lumbar disc extrusion. *Am J Roentg* June 1989;152(6):1277-1289.
57. Sterling M, Jull G, Wright A. Cervical mobilisation: concurrent effects on pain, sympathetic nervous system activity and motor activity. *Man Ther.* 2001;6(2):72-81
58. Teodorczyk-Injeyan JA, Injeyan HS, Ruegg R. Spinal manipulative therapy reduces inflammatory cytokines but not substance P production in normal subjects. *J.Manipulative Physiol Ther.* 2006;29:14–21.
59. Vernon HT, Dhimi MS, Howley TP, Annett R. Spinal manipulation and beta-endorphin: a controlled study of the effect of a spinal manipulation on plasma beta-endorphin levels in normal males. *J Manipulative Physiol Ther.* 1986;9:115–123.
60. Vicenzino B, Collins D, Benson H, Wright A. An investigation of the interrelationship between manipulative therapy-induced hypoalgesia and sympathoexcitation. *J Manipulative Physiol Ther.* 1998;21:448–453.
61. Wright, A. Hypoalgesia post-manipulative therapy: a review of a potential neurophysiological mechanism. *Man Ther.* 1995;1:11–16.
62. Pierau FK, Fellmer G, Taylor DC. Somato-visceral convergence in cat dorsal root ganglion neurons demonstrated by double labelling with fluorescent tracers. *Brain Res.* 1984;321(1):63-70.
63. Kimura A, Sato A. Somatic regulation of autonomic functions in anesthetized animals – neural mechanisms of physical therapy including acupuncture. *Jpn J Vet Res.* 1997;45(3):137-145.
64. Sato A, Sato Y, Schmidt RF. The impact of somatosensory input on autonomic functions.. In: *Reviews of Physiology Biochemistry and Pharmacology.* Blaustein MP et al Eds. Springer-Verlag, Berlin. 1997;v130.
65. Sato A. Neural mechanisms of somatosensory regulation of catecholamine secretion from the adrenal gland. *Adv Biophys.* 1987;23:39-80.
66. Haavik-Taylor H, Murphy B. Altered central integration of dual somatosensory input after cervical spine manipulation. 2008;33(3):178-188.
67. Haavik-Taylor H, Murphy B. The effects of spinal manipulation on central integration of dual somatosensory input observed after motor training: a crossover study. *J Manipulative Physiol Ther.* 2010;33(4):261-272.
68. Haavik H, Niazi IK, Holt K, Murphy B. Effects of 12 weeks of chiropractic care on central integration of dual somatosensory input in chronic pain patients: a preliminary study. *J Manipulative Physiol Ther.* 2017;40(3):127-138
69. Pickar JG, Bolton PS. Spinal manipulative therapy and somatosensory activation. *J Electrophys Kinesiol.* 2012;22(5):785-794
70. Gu H. [Observations on somatosensory evoked potential changes before and after traditional manipulation for cervical radiculospodylosis.] *Zhong Xi Yi Jie He Za Zhi* 1986;6(12):734-736. (Chinese) PMID 2951026
71. Haker E, Egekvist H., Bjerring P. Effect of sensory stimulation (acupuncture) on sympathetic and parasympathetic activities in healthy subjects. *J. Auton. Nerv. Syst.* 2000;79: 52–59. 10.1016/S0165-1838(99)00090-9.
72. Desmedt JE. Critical neuromonitoring at spinal and brainstem levels by somatosensory evoked potentials. *Cent Nerv Syst Traume.* 1985;2(3):169-186.
73. Heiskari M, Tolonen U, Kovala T et al. Somatosensory evoked potentials to posterior tibial nerve stimulation in cervical radiculopathy and radiculomyelopathy. *Neuro Orthop* 1988;5(2):78-82.
74. Capria MP. Somatosensory neurological evaluation of chiropractic manipulation. *J Chiro Research & Clin Investigation* 1990;6(3):56-58.
75. Bialosky JE, Bishop MD, Price DD, Robinson ME, George SZ. The mechanisms of manual therapy in the treatment of musculoskeletal pain: a comprehensive model. *Man Ther* 2009;14(5):531-538.
76. Daligadu J, Haavik H, Yelder PC, Baarbé J, Murphy B. Alterations in cortical and cerebellar motor processing in subclinical neck pain patients following spinal manipulation. *J Manipulative Physiol Ther.* 2013;36(8):527-537
77. Haavik H, Murphy B. Subclinical neck pain and the effects of cervical manipulation on elbow joint position sense. *J Manipulative Physiol Ther.* 2011;34(2):88-97
78. Farid B, Yelder P, Holmes M, Haavik H, Murphy BA. Association of subclinical neck pain with altered multisensory integration at baseline and 4-week follow-up relative to asymptomatic controls. *J Manipulative Physiol Ther.* 2018;41(2):81-81.

79. Kelly D, Murphy B, Backhouse D. Use of a mental rotation reaction-time paradigm to measure the effects of upper cervical adjustments on cortical processing: a pilot study. *J Manipulative Physiol Ther* 2000;23:246-251.
80. Luke M. Subclinical neck pain impairs cognitive ability which can be improved by chiropractic treatment: a four week longitudinal study with a healthy control group comparison. *Internat'l Conference Pain Research and Management*. (See also *J Pain Relief*. DOI:10.4172/2167-0846.C1.012)
81. Romero E. The cerebellum and central dysdiadokinesia. *Dynamic Chiropr* 2009;27(7). <https://www.dynamicchiropractic.com/mpacms/dc/article.php?id=53708>
82. Sergeant A. Chiropractic management of a patient with cerebellar atrophy. [Case Report]. *Chiropr J Aust* 2017;45:175-183.
83. O'Sullivan PB, Burnett A, Floyd AN, et al. Lumbar repositioning deficit in a specific low back pain population. *Spine* 2003;28(1):1074-79
84. Reed WR, Pickar JG, Sozio RS, Long CR, Effect of spinal manipulation thrust magnitude on trunk mechanical activation thresholds of lateral thalamic neurons. *J Manipulative Physiol* 2014;37(5):277-286. Earlier
85. Reed WR, Cranston JT, Onifer SM, Sozio RS. Decreased spontaneous activity and altered evoked nociceptive response of rat thalamic submedial neurons to lumbar vertebra thrust. *Exp Brain Res* 2017;235(9):2883-2892
86. Yang SW, Follett KA, Piper JG, Ness TJ. The effect of morphine on responses of mediodorsal thalamic nuclei and nucleus submedialis neurons to colorectal distension in the rat. *Brain res* 1998;779(1-2):41-52
87. Liu X. [The modulation of cerebral cortex and subcortical nuclei on NRM and other role in acupuncture analgesia]. *Zhen Ci Yan Jiu* 1996;21(1):4-11 (English extract)
88. Vaganian LG, Bagdasarian RA, Karapetian DS. [Neuronal organization of somatovisceral afferent systems of the centromedian area of the thalamus]. *Fiziol Zh SSSR Im I Sechanova*. 1985;71(1):72-79. (Abstract only)
89. Boal RW, Gillette RG. Central neuronal plasticity, low back pain and spinal manipulative therapy. *J Manipulative Physiol Ther*. 2004;27(5):314-326.
90. Gay CW. Immediate changes after manual therapy in resting state functional connectivity as measured by functional magnetic resonance imaging in participants with induced low back pain. *J Manipulative Physiol Ther*. 2014;37(9):614-627.
91. Cortelli P, Giannini G, Favoni V, Cevoli S, Pierangeli G. Nociception and the autonomic nervous system. *Neurol Sci* 2013;34 (Suppl 1):S41-46
92. Hamunen K, Kontinen V, Hakala E, Talke P, Paloheimo M, Kalso E. Effect of pain on autonomic nervous system indices derived from photoplethysmography in health volunteers. *Brit J Anaesth*. 2012;108(5):838-844.
93. Schlereth T, Birklein F. The sympathetic nervous system and pain. *Neuro Molec Med* 2008;10(3):141-147
94. Diatchenko L, Fillingim RB, Smith SB, Maixner W. The phenotypic and genetic signatures of common musculoskeletal pain conditions. *Nat Rev Rheumatol* 2013;9(6):340-350
95. Crofford L J. Chronic pain: where the body meets the brain. *Trans Am Clin Climatol Assoc*. 2015;126:167-183.
96. Appelhans BM, Lueken LJ. Heart rate variability and pain: associations of two interrelated homeostatic processes. *Biol Psychol* 2008;77(2):174-182.
97. Koenig J, Jrczok MN, Ellis RJ, Hillecke TK, Thayer JF. Heart rate variability and experimentally induced pain in healthy adults: a systematic review. *Eur J Pain* 2014;18(3):301-314
98. Woolf CJ. Nociceptors – noxious stimulus detectors. *Neuron* 2007;55(3):353-364.
99. Budgell B. Autonomic responses to spinal pain. *Rigakaryoha Kagaku* 2000;15(3):81-87.
100. Benarroch EE. Pain-autonomic interactions: a selective review. *Clin Autonomic Res* 2001;11:343-349.
101. Pace MC, Mazhariello L, Passavanti MB, Sansone P, Barbarisi M, Aurilio C. Neurology of pain. *J Cell Physiol* 2006;209(1):8-12
102. Apkarian AV, Bushnell MC, Treede RD, Zubieta JK. Human brain mechanisms of pain perception and regulation in health and disease. *Eur J Pain* 2005;9(4):463-484.
103. Mehl-Madrona L, Mainguy B. Neurobiology of pain. <http://www.mehl-madrona.com/Neurobiology%20of%20Pain%20for%20APA%202012.pdf>
104. Sillevs R, Cleland J. Immediate effects of the audible pop from a thoracic spine thrust manipulation on the autonomic nervous system and pain. A secondary analysis of a randomised clinical trial. *J Manipulative Physiol Ther* 2011;34(1):37-45
105. Husain JM, LaRoche M, Keosaian J, et al. Reasons for opioid discontinuation and unintended consequences following opioid discontinuation within the TOPCARE trial. *Pain Med* 2018; doi: 10.1093/pm/pny124.
106. Scherbaum N, Bonnet U. [Neurobiology of opioid dependence.] *Schmerz* 2018; doi: 10.1007/s00482-018-0304-8. (Abstract)
107. Vigotsky AD, Bruhns RP. The neurophysiological response to manual therapy and its analgesic implications: a narrative review. (Thesis). Arizona State University. 2015, June 5. <https://peerj.com/preprints/996.pdf>.
108. Srbely J, Vernon H, Lee D, Polgar M. Immediate effects of spinal manipulative therapy on regional antinociceptive effects in myofascial tissues in healthy young adults. *J Manip Physiol Ther*. 2013;36(6):333-341.

- 109.Burton AR, Fazalbhoy A, Macefield VG. Sympathetic responses to noxious stimulation of muscle and skin. *Front Neurol*.2016;7:109: 10.3389/fneur.2016.00109.
- 110.Hawk C, Cambron J. Chiropractic care for older adults: effects on balance, dizziness, and chronic pain. *J Manipulative Physiol Ther* 2009;32(6):431-437.
- 111.Holt KR, Noone PL, Short K, Haavik H. Fall risk profile and quality-of-life status of older chiropractic patients. *J Manipulative Physiol Ther* 2011;34(2):78-87
- 112.Holt KR, Haavik H, Elley CR. The effects of manual therapy on balance and falls” a systematic review. *J Manipulative Physiol Ther* 2012;35(3):227-234
- 113.Norre ME. Neurophysiology of vertigo with special reference to cervical vertigo. A review. *ACTA Belgica Medica Physica* 1986;9(3)183-194.
- 114.Polus BJ. Muscle spindles and spinal proprioception. In: Haldeman S. *Principles and Practice of Chiropractic*. 3rd edn. New York: McGraw-Hill Education; 249-288.
- 115.Pickar JG, Bolton PS. Spinal manipulative therapy and somatosensory activation. *J Electromyogr Kinesiol* 2012;22(5):785-794
- 116.Purves D, Augustine GJ, Fitzpatrick D, et al. eds. *Mechanoreceptors specialised for proprioception*. Neuroscience 2nd edn. Sunderland: Sinauer Associates. 2001. <https://www.ncbi.nlm.nih.gov/books/NBK10812/>
- 117.Swinkels A. Spinal position sense is independent of the magnitude of movement. *Spine* 2000;25(1):98-104
- 118.Swinkels A, Dolan P. Spinal position sense and disease progression in ankylosing spondylitis: a longitudinal study. *Spine* 2004;29(11):1240-1245.
- 119.Gong W. Effects of cervical joint manipulation on joint position sense of normal adults. *J Phys Ther Sci*. 2013;25(6):721-723.
- 120.Grgić V. [Cervicogenic proprioceptive vertigo: etiopathogenesis, clinical manifestations, diagnosis and therapy with special emphasis on manual therapy. *Lijec Vjesn*. 2006;128(9-10):288-295.
- 121.Korr IM. Proprioceptors and somatic dysfunction. *J Am Osteopath Assoc*. 1975;74:638–650.
- 122.Learman KE, Myers JB, Lephart SM, et al. Effects of spinal manipulation on trunk proprioception in subjects with chronic low back pain during symptom remission. *J Manipulative Physiol Ther*. 2009;32(2):118-126.
- 123.Marshall LD. The effect of cervical spine manipulation on elbow joint position sense in patients with acute cervical facet syndrome. Thesis. University of Johannesburg. <https://ujcontent.uj.ac.za/vital/access/manager/Repository/uj:7555/CONTENT1>
- 124.Miranda F, Faccini D, Manfio F. Influence of cervical spine manipulation on neck joint position sense error in patients with chronic neck pain. *Man Ther Posturo Rehab J*. 2016; 1-5. <https://doi.org/10.17784/mtprehabJournal.2015.13.283>
- 125.O’Sullivan PB, Burnett A, Floyd AN, et al. Lumbar repositioning deficit in a specific low back pain population. *Spine* 2003;28(1):1074-1079
- 126.Slosberg M. Effects of altered afferent articular input on sensation, proprioception, muscle tone and sympathetic reflex responses. *J Manipulative Physiol Ther*. 1988;11:400-08.
- 127.Stanton TR, Leake HB, Chalmers KJ, Moseley GL. Evidence of impaired proprioception, idiopathic neck pain: systematic review and meta-analysis. *Phys Ther* 2016;96(6):868-887
- 128.Zimny ML. Mechanoreceptors in articular tissues. *Am J Anat* 1988;182(1):16-32.
- 129.Harris W, Wagnon RJ. The effects of chiropractic adjustments on distal skin temperature. *J Manipulative Physiol Ther*; 1987;10(2):57-60
- 130.Korr IM, Wright HM, Chace JA. Cutaneous patterns of sympathetic activity in clinical abnormalities of the musculoskeletal system. In: Peterson B, ed. *The collected papers of Irvin M Korr*. Newark. American Academy of Osteopathy. 1979;66-72.
- 131.Thomas PE. Sympathetic activity in facilitated segments: sudomotor studies. *J Am Osteop Assoc*. 1955;55;5-8.
- 132.Vicenzino, B., Collins, D., Wright, T. Sudomotor changes induced by neural mobilisation techniques in asymptomatic subjects. *J Man Manip Ther*. 1994;2:66–74.
- 133.Zwaafink KG. Neurovegetative reactions of spinal manipulations and mobilisations in manual therapy, chiropractic and osteopathic medicine. Master’s Thesis. Tyrol. Univ Applied Sciences. July 2016.
- 134.Cunningham BM, Grimm DR, Burke JR. Autonomic nervous system function among individuals with acute musculoskeletal injury. *J Manipulative Physiol Ther* 2005;28(1):44-51.
- 135.Sato A, Sato Y, Schmidt RF. (64) p 259.