# Neurodynamics of vertebrogenic somatosensory activation and Autonomic Reflexes - a review:

Part 7 The cervicogenic factor

Peter Rome and John Waterhouse

Abstract: Articular activation of sensory nerves should not be seen as being limited to nociceptive receptors as other neural sensory pathways are also vulnerable to noxious stimulation.

Indexing terms: Vertebral subluxation; cervicogenic; Somatosensory; Autonomic nervous system.

## Introduction

### 'Cervical dysfunction is the most common cause of neck pain.' (1)

D ue to its anatomical proximity to the brainstem, a disturbed cervical vertebra appears to have the potential to adversely influence a number of neural structures and physiological functions. As such, this neurophysiological impact may contribute a profound neurospinal component upon the autonomic nervous system - especially the brainstem nuclei of the parasympathetic cranial nerves. (See also Cranial Nerves Part VIII) (2, 3, 4, 5, 6)

As nociception is a neurological phenomenon, it qualifies as a somatosensory response. Cervicogenic headaches are another example. Both examples of neural reflex disturbance indicate that cervical subluxations or dysfunction are more than just osseous disturbance. (7, 8)

The cervical spine has featured strongly in studies on spinal influence upon the ANS. Three examples of these are cervicogenic headaches, cervical vertigo, and the cervical vertebral impact on particular cranial nerves. In whiplash injuries, a number of authors report classic examples of the cervicogenic pathophysiological neural symptoms of sensory, motor, and autonomic dysfunction associated with mechanical disruption of the cervical spine. Uchida and Budgell suggest that monitoring of the ciliospinal reflex as a possible diagnostic aid in assessing the

neural status of the cervical sympathetic trunk. (9)

Although they acknowledge medical controversy regarding cervicogenic headaches, Antonaci and colleagues recognise involvement of the nucleau cordalis and that *'Neuroimaging and* 

... cervical spine subluxations can affect 'central corticomotor facilitory and inhibitory neural processing and cortical motor control' involving the upper extremities.'





*kinematic analysis of neck motion may aid in diagnosing difficult CEH*' (Cervicogenic Headache). Four years later Vincent stated that the '*Cervicogenic headache (CEH) is a well-recognized syndrome,*' *and 'a relatively common syndrome*.' (10, 11, 12)

It is noted that while there is increasing literature and evidence of the condition, there seems an implied reluctance to cite refereed papers from chiropractic or osteopathic sources – the principal proponents in the field. A medical alternative which also recognises the cervicogenic nature of headaches would be interventionist measures such as nerve blocks. Although Edmeads suggested that nerve blocks are of temporary benefit and should not be administered frequently. (13, 14, 15)

Headaches, migraines, cardiac and respiratory changes feature as being regularly associated in cervicogenic-related conditions. Using Wistar rat subjects, a potentially pertinent paper by Edwards et al in 2015 outlined potential adverse effects of a disrupted cervical spine. This suggests a physiologically functional (mechanical) normalised cervical spine would tend not to experience those symptoms. The authors found that noxious sensory stimulation from whiplash involving the C2 spinal nerve, can precede cervical dystonia. Further, that this could influence the autonomic pathways resulting in such conditions as cardiovascular and respiratory abnormalities, dysphagia, and *'eye position'* disturbances. (16, 17)

Also in relation to cervical dysfunction headaches, Braaf and Rosner suggested that 90% of headaches are cervicogenic in nature. They published seven studies on cervical disturbances over the period 1953 to 1975. (18, 19, 20, 21, 22, 23, 25, 26)

In further animal experiments, Edwards et al established that the *intermedius nucleus* of the medulla (InM) receives input from the cervical region leading to autonomic responses. It is also associated with the coordination of tongue movements. (16)

Additional research by Edwards and colleagues investigated neurological sensory afferent links between the upper cervical spine (namely the C2 nerve), postural control, mouth and eye movement, as well as respiratory and cardiovascular symptoms. They also identified a proprioceptive association through a group of brainstem nuclei. In particular the research noted that stimulation of the intermedius nucleus (InM) 'mimicked the response of second cervical nerve stimulation'. Their findings tend to corroborate the link between the upper cervical sensory afferents and the autonomic nervous system control through the InM. They also recognise the potential for dysphagia and a form of cardiorespiratory dysfunction which may be associated with a disturbed cervical spine, and cervical dystonia resulting from whiplash injuries. (27)

Vaňásková et al also noted that in cervicogenic dysphasia the swallowing reflex may involve five of the cranial nerves: *Trigeminal* (V), *Facial* (VII), *Glossopharyngeal* (IX), *Vagus* (X), and the *Hypoglossal* (XII). (28)

In extensive research involving cervicogenic disorders, Haavik-Taylor and colleagues have demonstrated that cervical spine subluxations can affect '*central corticomotor facilitory and inhibitory neural processing and cortical motor control*' involving the upper extremities. This central processing could also be indicative of the wider implications of neural effects from mechanical vertebral disturbances, VSCs. (29)

The ENT surgeons Franz and colleagues suggested that cases of cervicogenic oto-ocular syndrome could be due to vertebral disturbance (functional disorders) associated with such signs and symptoms as, mydriasis, sensorineural hearing loss, vertigo, tinnitus, and *Eustachian tube* dysfunction. In the study, patients' ANS function was monitored by observing pupil size. The cervical spine was assessed by a physical therapy examination of facet mobility. (31)

The neck-tongue syndrome may also be seen as being cervicogenic in nature. It is characterised by associated neck or sub-occipital pain. As headaches usually accompany this condition, the syndrome is now classified in the *International Classification of Headache Disorders* (ICHD-3) as item code 13.5, *Neck-tongue syndrome*. In an indication of a spine-related biomechanical factor of its onset, the ICHD notes that pain associated with the syndrome is precipitated by a '*sudden turning of the head*'. Effective chiropractic management of the condition has been reported. (32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45)

In further research, Bolton and colleagues noted a vestibulo-sympathetic correlation with posture involving dorsal neck muscles, cervical afferents (C2, C3) with cardiovascular and respiratory responses in cat subjects. (46, 47)

Others studies indicate that birth injuries to an infant's neck may be a precursor to a range of cervicogenic conditions, including infantile colic. The physicality and the infant's vulnerability during the birth process are considered to render them particularly susceptible to cervical injury. Naturally, conventional chiropractors adapt light adjusting techniques on such patients. (49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 61, 63, 64, 65, 66, 67, 68, 69)

Rostocki stated that 'Many infants suffer neck injury during birth which can lead to lifelong health problems.......These injuries can be caused organically or through medical error and may affect any structure or system in the newborn infant's body. The cervical spinal column is very delicate in an infant and is often the site of trauma from a birth injury.' (70) He also stated that 'Chiropractic for neck pain is perhaps the most common and popular complementary healing art used by patients the world over.' (71)

Interpreting newborns' spine-related symptoms is a regular paediatric procedure for most chiropractors. (72, 73, 74, 75) Pizzolorusso and colleagues also found a rather extensive range of subluxation (osteopathic somatic dysfunctions) in 155 infant paediatric patients. (76)

Durand and Daniels investigated the compression of the C4 nerve root. They noted such symptoms as phrenic nerve palsy, shortness of breath, chest pain, and an elevated hemidiaphragm. While this is an example of a marked physical state, it is suggested that lesser segmental disturbances, irritation, or neural sensory activation of the C4 may also have the potential to lead to symptoms related to the structures involved, particularly when of a chronic nature. (77)

# Conclusion

The listed variety of cervicogenic disorders would suggest that comparative studies of prevalence and therapeutic efficacy are warranted



John D Waterhouse DC, FACC Private practice, Melbourne

*Cite:* Rome P. Waterhouse JD. Neurodynamics of vertebrogenic somatosensory activation and Autonomic Reflexes - a review: Part 7 The Cervicogenic Factor. Asia-Pacific Chiropr J. 2021;1.4. URL apcj.net/papers-issue-2-4/ #RomeWaterhousePart7Cervicogenic

### References

- 1. Health Engine. https://healthengine.com.au/info/cervical-dysfunction Jan 1, 2004 (Extracted 17.1.2021.)
- Bigland MJ, Budgell BS, Bolton PS. The influence of cervical spinal cord compression and vertebral displacement on somatosympathetic reflexes in the rat. Spine J. 2015 Jun 1;15(6):1310-1317. DOI 10.1016/j.spinee.2013.08.013. Epub 2013 Oct 28.
- 3. Cramer GD. The cervical region. Chapter 5. In: Cramer GD, Darby SA. Basic and clinical anatomy of the spine, spinal cord, and ANS. St Louis,. Mosby. 1995:144-5.
- Foreman SM. Nervous system trauma. In: Foreman SM, Croft AC. (eds). Whiplash injuries. The cervical acceleration/deceleration syndrome. Third edn. Philadelphia: Lippincott Williams & Wilkins. 2002:429-52.
- 5. Gatterman MI. Whiplash: a patient-centered approach to management. St Louis: Elsevier Mosby. 2012.
- 6. Bogduk N. Ther anatomy and physiology of whiplash. Clin Biomech 1986;1:92-101.
- 7. Passadore M, Roatta S. Influence of sympathetic nervous system on sensorimotor function: whiplash associated disorders (WAD as a model. Eur J Appl Physiol. 2006;98:423-49.
- 8. Passadore M,Deriu F, Roatta S, Grassi C, Micieli G. Effects of cervical sympathetic nerve stimulation on the cerebral microcirculation: possible clinical implications. Acta Neurobiol Exp. 1996;56:117-27.
- 9. Uchida S, Budgell B. Somato-autonomic reflex. Encyclopaedia Neurosci. 2009;3767-3770 DOI 10.1007/978-3-540-29678-2\_5500.
- 10. Antonaci F, Bono G, Chimento P. Diagnosing cervicogenic headache. J Headache Pain. 2006;7(3):145-8.
- 11. Vincent MB. Cervicogenic headache: a review comparison with migraine, tension-type headache, and whiplash. Curr Pain Headache Rep. 2010;14(3):238-43.
- 12. Vincent MB. Headache and neck. Curr Pain Headache Rep. 2011;15(4):324-31.
- 13. Werner J, Becker WJ. Cervicogenic headache: evidence that the neck is a pain generator. Headache. 2010;50(4):699-705.
- 14. Yi X, Cook AJ, Hamill-Ruth RJ, Rowlingson JC. Cervicogenic headache in patients with presumed migraine: missed diagnosis or misdiagnosis? J Pain. 2005;6(10):700-3.
- 15. Edmeads J. Cervicogenic headache. Pain Res Man. 1996;1(2):119-22.
- Edwards IJ, Lall VK, Paton JF, et al. Neck muscle afferents influence oromotor and cardiorespiratory brainstem neural circuits. Brain Structure Function 2015;220(3):1421-36.
- 17. Moore C, Adams J, Leaver A, Lauche R, Sibbritt D. The treatment of migraine within chiropractic: analysis of a nationally representative survey of 1869 chiropractors. BMC Compl Altern Med 2017;17:519. https://doi.org/10.1186/s12906-017-2026-2033.
- 18. Butler S. Headaches a real pain-in-the-neck. Sydney Telegraph. August 27th 1970 pp7. (Citing Drs Braaf & Rosner).
- 19. Braaf MM, Rosner S. The treatment of headaches. New York State J Med. Mar 15, 1953;53:687-93.
- 20. Braaf MM, Rosner S. Symptomatology and treatment of injuries of the neck. New York State J Med. Jan 15, 1955;55:237-42.
- 21. Braaf MM, Rosner S. Whiplash injury of the neck: Symptoms, diagnosis, treatment, and prognosis. New York State J Med. 1958;58:1501-7.
- 22. Braaf MM, Rosner S. Chronic headache. A study of over 2000 cases. New York State J Med. Dec 15, 1960;3987-95.
- 23. Braaf MM, Rosner S. Meniere-like syndrome following whiplash injury of the neck. J Trauma. 1962;2:494-501.
- 24. Braaf MM, Rosner S. Headaches in children. Headache. 1970;9:207-10.
- 25. Braaf MM, Rosner S. Trauma of cervical spine as cause of chronic headache. J Trauma. 1975;15(5):441-6.
- 26. Çoban G, Çöven I, Çifçi BE, Yildirim E, Yazici AC. The importance of craniovertebral and cervicomedullary angles in cervicogenic headache. Diagn Interv Radiol. 2014;20(2):172-77.
- 27. Edwards IJ, Deuchars SA, Duechars J, The intermedius nucleus of the medulla: a potential site for the integration of cervical information and the generation of autonomic responses. J Chem Neuroanat 2009;38(3):166-75.
- Vaňásková E, Hep A, Vižď a J, Tosnerová V. Swallowing disorders related to vertebrogenic dysfunction. Ceska a Slovenska Neurologie a Neurochirurgie 2007;70(6):692-696. In: Brzozowski T (Ed.). New advances in the basic and clinical gastroenterology., ISBN: 978-953-51-0521-3; 2012. Rijeka, Croatia; In: Tech. 2012 p175-184 http://www.intechopen.com/books/new-advances-in-the-basic-andclinicalgastroenterology/swallowing-disorders-related-to-vertebrogenic-dysfunctions.
- 29. Haavik-Taylor H Murphy B. Cervical spine manipulation alters sensorimotor integration evoked potential study. Neurophysiol. 2007;118(2):391-402.
- Franz B, Altidis P, Altidis B, Collis-Brown G. The cervicogenic otoocular syndrome: a suspected forerunner of Ménère's disease. Int Tinnitus J. 1999;5(2):125-139. http://www.tinnitusjournal.com/articles/the-cervicogenic-otoocular-syndromea-suspected-forerunner-ofmenieres-disease.pdf.

- 31. Franz B, Collis-Brown G, Altidis P, Altidis B, Cummings N. Cervical trauma and tinnitus. Int Tinnitus J 1998;4(1):31-3.
- 32. IHS Classification ICHD-3. Painful lesions of the cranial nerves and other facial pain. https://www.ichd-3.org/13-painful-cranialneuropathies-and-other-facial-pains/13-5-optic-neuritis/.
- 33. Borody C. Neck-tongue syndrome. J Manipulative Physiol Thor. 2004;27(5):e8.
- 34. Buroon M, Pero J. Resolution of glossopharyngeal neuralgia & spastic dystonia following chiropractic care to reduce upper cervical vertebral subluxation: A case study [case report. J Upper Cervical Chiropr Res. 2014:1:7-13.
- Cassidy JD, Diakow PRP, De Korompay VL, Munkacsi I, Yong-Hing K. Treatment of neck-tongue syndrome by spinal manipulation. Pain Clinic. 1986;1(1):41-46. https://pdfs.semanticscholar.org/3cf6/f72448957512d1548c97701583cd02f233f3.pdf.
- 36. Gelfand AA, Johnson H, Lenaerts ME, et al. Neck-tongue syndrome: a systemic review. Cephalalgia. 2018;38(2):374-382
- 37. Hankey, GJ. Neck-tongue syndrome on sudden neck rotation. Aust N Z J Med. 1988; 18: 181.
- Lavigne V. A narrative review and case report. Frenotomy procedure in neonate with tongue-tie. J Clin Chiropr Pediatr. 2012;13(2):1025-31.
- 39. Lois I. The cervicolingual syndrome. Man Med. 1987; 3: 63-6.
- 40. Niethamer L, Myers R. Manual therapy and exercise for a patient with neck-tongue syndrome a case report. J Orthop Sports Phys Ther. 2016;46(3):217-224.DOI 10.2519/jospt.2016.6195. (Ed. note: this is designated as C2).
- 41. Sjaastad O, Bakketeig LS. Neck-tongue syndrome and related (?) conditions. Cephalalgia. 2006;26(3):233-40.
- 42. Lance JW, Anthony M. Neck-tongue syndrome on sudden turning of the head. J Neurol Neurosurg Psychiatry. 1980;43:97-101.
- 43. Terrett AGJ. Neck-tongue syndrome and spinal manipulative therapy. In: H Vernon (Ed.) Upper cervical syndrome: chiropractic diagnosis and treatment. Williams & Wilkins, Baltimore; 1988: 223–9.
- 44. McGuiness J., Vicenzino, B., Wright, A. Influence of a cervical mobilization technique on respiratory and cardiovascular function. Man Ther. 1997;2:216–20.
- 45. Vicenzino B, Cartwright T., Collins D, Wright, A. Cardiovascular and respiratory changes produced by lateral glide mobilization of the cervical spine. Man Ther. 1998;3:67–71.
- 46. Bolton PS, Kerman JA, Woodring SF, Yates BJ. Influences of neck afferents on sympathetic and respiratory nerve activity. Brain Res Bull. 1998;47(5):413-19.
- 47. Bolton PS, Ray CA. Neck afferent involvement in cardiovascular control during movement. Brain Res Bul .2000;53(1):45-9.
- 48. Bagagiolo D, Didio A, Sbarbaro M, Peiolo CG, Borro T, Farina D. Osteopathic manipulative treatment in pediatric and neonatal patients and disorders: clinical considerations and updated review of the existing literature. Am J Perinatalol. 2016;33:1050-4.
- 49. Biedermann H. Kinematic imbalances due to suboccipital strain in newborns. J Man Med. 1992;6:151-6.
- 50. Biedermann H. The KISS syndrome: symptoms and signs. In: Biedermann H (ed). Manual therapy in children. New York: Churchill Livingstone; 2004;285-302.
- 51. Caird MS, Reddy S, Ganley TJ, Drummond DS. Cervical spine fracture-dislocation birth injury: prevention, recognition, and implications for the orthopaedic surgeon. J Pediatr Orthop. 2005;25(4):484-86.
- 52. Cerritelli F, Martelli M Barlafante G. Introducing an osteopathic approach into neonatology ward: the NE-O model. Chiropr Man Ther. 2014;22:18.
- 53. Cintaretu G. A review of osteopathic treatments for paediatric conditions. Jan 7, 2016 https://www.numss.com/Thesis/ Gigel%20Cintaretu%20NUMSS%20Thesis.pdf
- 54. Collins K, Alcantara J, Holt K. resolution of breast feeding and gastrointestinal complaints in infant twins with plagiocephaly and scaphocephaly following birth trauma. J Pediatr Maternal Family Health 2015;1:34-41.
- 55. Fallon JM, Fysh PN. Chiropractic care of the newborn with congenital torticollis. J Clinical Chiropr Pediatr. 1997;2(1):116-21.
- 56. Siegenthaler MH. Chiropractic management of infantile torticollis with associated abnormal fixation of one eye: a case report. J Chiropr Med. 2015;14(1):51-6.
- 57. Gottlieb M. Neglected spinal cord, brain stem, and musculoskeletal injuries stemming from birth trauma: a review of the literature. J Manipulative Physiol Ther. 1993;16:537-43.
- 58. Guliana K, Rubin D. Improvements in developmental delay, colic and GERD in a child undergoing chiropractic care: a case report and review of literature. J Pediatr, Maternal Family Health.2012;1:1-4.
- 59. Hughes CA, Harley EH, Milmoe G, Bala R, Martorella A. Birth trauma in the head and neck. Arch Otolaryngol Head Neck Surg. 1999;125(2):193-9.
- 60. Kraus LL. Case study: birth trauma results in colic. Chiropr Pediatr. 1995;2:10-11.
- Neally R, Alcantara J. Resolution of failure to thrive in a 4-month-old infant following adjustment of subluxation: case and selective review of the literature. J Pediatr Maternal Family Health. 2015;4:149-53.

- 62. Pizzolorusso G, Cerritelli F, D'Orazio M, et al. Osteopathic evaluation of somatic dysfunction and craniosacral strain pattern among preterm and term newborns. J Am Osteopath Assoc. 2013;113(6):462-7.
- 63. Pizzolorusso G, Turi P, Barlafante G. et al. Effect of osteopathic manipulative treatment on gastrointestinal function and length of stay of preterm infants: an exploratory study. Chiropr Man Ther. 2011;19:15.
- 64. Ritzmann D. Birthing interventions and the newborn. In Biedermann H. (50 p 85-4).
- 65. Rome P, Waterhouse J, Maginness G, Ebrall P. Medical management of infantile colic with spinal manipulation: a narrative review of European medical literature. J Contemporary Chiropr. 2019;2:54-69.
- 66. Rome P. Medical management of paediatric and non-musculoskeletal conditions by spinal manipulation. Chiropr J Aust. 2013;43(4):131-8.
- 67. Spear D, Alcantara J. Resulution of birth trauma sequelae following adjustment of vertebral subluxations in an infant. J Pediatr Maternal Family Health. 2016;1:28-31.
- 68. Towbin A, Latent spinal cord and brain stem injury in newborn infants. Develop Med Chil Neurol. 1969;11:54-68.
- 69. Waddington EL, Snider KT, Lockwood MD, Pazdernik VK. Incidence of somatic dysfunction in healthy newborns. J Am Osteop Assoc. 2015;115(11)654-65.
- 70. Rostocki A. Neck injury during birth. https://www.neck-pain-treatment.org/neck-injury-during-birth.html.
- 71. Rostocki A. Neck pain treatment. https://www.neck-pain-treatment.org/neck-pain-treatment.html.
- 72. Davies N. Chiropractic paediatrics: a clinical handbook. Edinburgh: Churchill Livingstone. 2000.
- 73. Anrig C, Plaugher G. Pediatric chiropractic. Philadelphia. Lippincott Williams Wilkins. 2012.
- 74. Biedermann H (ed). Manual therapy in children. New York: Churchill Livingstone. 2004.
- 75. Painter FM. Chiropractic pediatrics. https://chiro.org/pediatrics/.
- 76. Pizzolorusso G, Cerritelli F, D'Orazio M, et al. Osteopathic evaluation of somatic dysfunction and craniosacral strain pattern among preterm and term newborns. J Am Osteop Assoc. 2013;115(6):462-7.
- 77. Durand WM, Daniels AH. C4 root compression leads to phrenic nerve palsy with shortness of breath, chest pain, and elevated hemidiaphragm. Lancet. 2020;396(10257):1101. DOI 10.1016/S0140-6736(20)32027-4.