

# Neurodynamics of vertebrogenic somatosensory activation and Autonomic Reflexes - a review:

## Part 5 Diversity in Vertebral Subluxations

Peter Rome and John Waterhouse

**Abstract:** Elements of the subluxation model are well known but other aspects are still emerging. Some remain in a contextual sense. This review presents a synopsis of currently available hypotheses on the types of segmental disturbances, the need for specificity, research involving animal subjects, and why the focussed adjustment to normalise dysfunctional segments is a significant part of the chiropractic model.

**Indexing terms:** Vertebral subluxation; Neurophysiology; Somatosensory; Autonomic nervous system

### Introduction

**A**s distinct from major tissue damage or pathology, the somatovisceral model discussed here is potentially a clinical pathophysiological contributor involving noxious or aberrant neural afference of somatic origin.

These are known to activate autonomic nervous system – somato-autonomic reflexes. The vertebral subluxation complex (VSC) - especially the disrupted vertebral articular elements, would appear to be a primary initiator of the noxious neural reflex arcs. These can be associated with somato-autonomic, somatovisceral and somatosympathetic, somatosomatic, somata-parasympathetic reflexes, with hypo or hyperreflexia pathophysiology. (1, 2, 3, 4, 5 6, 7, 8, 9, 10, 11)

In order to clarify the complex role of disturbed vertebral physiology, the following definition of a subluxation is submitted.

A subluxation is an articular dysfunction, typically but not limited to the spine and pelvis, which is characterised by anatomical and neurophysiological signs and symptoms.

In a further clarification aimed at differentiating the advanced form of correcting subluxations, the following definition of an adjustment is submitted.

The spinal adjustment is defined here as:

subluxation

*...it would seem apparent that the subluxation is far more complex than a mere mechanical manipulative lesion.'*



The physical application of a highly developed and finely tuned advanced form of manual or instrument intervention directed to restoring joint and neural physiology in order to ameliorate associated signs and symptoms.

Other models of this lesion have also been proposed for a subluxation construct. Such a range of hypotheses would also suggest that there may be different forms of subluxations as well as being more than a simple mechanical displacement. Perhaps individual or various combinations of these models may be the somatic component activating somatovisceral reflexes. The range of subluxation models, or elements thereof reflect the complexity of the entity, they include:

- ▶ Biomechanical (12)
  - Intervertebral hypermobility (12)
  - Intervertebral hypomobility (13)
  - Intervertebral dyskinesia (13)
  - Intervertebral misalignment (12)
- ▶ Biomechanical fault (14)
- ▶ Dysafferentation Model (15)
- ▶ Facilitation Model (16)
- ▶ Functional neurology model (17)
- ▶ Malposition (18)
- ▶ Mechanical model (fixation, hypermobility) (19)
- ▶ Meniscoid entrapment (20)
- ▶ Neurobiologic model (21)
- ▶ Neurofasciogenic model (22)
- ▶ Neurological model (18)
- ▶ Nuclear fragmentation (19)
- ▶ Periarticular adhesions (23)
- ▶ Putative effects of the VSC (12)
  - Encroachment of intervertebral foramen or spinal canal
  - Altered afferent input from spinal and paraspinal tissues
  - Dentate ligament mediated cord distortion
- ▶ Sensory model (24)
- ▶ Somatosensory model (25)
- ▶ The fixation hypothesis (26)
- ▶ Trophic Model (18)
- ▶ Unified model for phases of VSC (27)
- ▶ Zygapophyseal joint sprain (28)

Senzon explored a detailed examination of the vertebral subluxation in a 10-part series with the models discussed in Part 9 in particular. (29) Other reviews have discussed the medical use of the subluxation model. (30, 31)

Despite reported amelioration following segmental adjustments of vertebral subluxations as recorded in the literature, (32, 33) this VSC entity has attracted controversy regarding its existence. (34) However, to the authors' knowledge, apart from superficial opinion and

unsubstantiated impressions and World Health Organisation recognition, (35) no formal evidential studies appear to have been produced to contradict it.

The target lesion for a specific adjustment must differ from other physically and functionally normal segments in order to have associated clinical signs and symptoms, and for them to respond to manipulative care. This would constitute cause and effect criteria which would justify conducting the corrective procedure of an adjustment.

In addition, there would be no valid reason in trying to manipulate a normally positioned and functioning asymptomatic vertebral segment. If the symptoms and signs are designated as vertebrogenic, then 'normal' functioning segments of the spine could hardly be held responsible. Unless presenting for ongoing supportive care where there may be no symptoms present, yet the clinician detects the presence of VSC's, much like a dentist may detect evidence of dental decay which is not yet symptomatic, a patient is unlikely to present themselves for care.

To question the legitimacy or effects of a *Somato-Autonomic Visceral Complex* (SAVC) hypothesis based merely on unsubstantiated opinion which lacks formal investigative research, demonstrates uninformed, unscientific, prejudicial bias. To do so is wilful ignorance of the available published evidence. Original scientifically researched evidence rejecting vertebral subluxation complex (VSC) concepts have not been forthcoming or seriously attempted.

As Stated by Sato et al, '*Thus the decrease in blood pressure and renal nerve activity during manipulation of the spine are thought to be supraspinal reflexes.*' (36) It would be more appropriate for those questioning the VSC to call for research to derive a definitive explanation as to why these segmental lesions seem directly related to patient symptoms, when a specific vertebral adjustment regularly alleviates those symptoms.

In 2011, Desmarais and colleagues demonstrated that thoracic spine manipulation may modulate somatosympathetic reflexes at a segmental level in health human subjects. Colloca, Keller et al. mechanically manipulated Merino sheep subjects with impulse instruments to study multi-axial vertebral segment motion. (37, 38, 39). Research of the science of subluxations and manipulation are noted both historically (40, 41), and in more recent reviews. (42, 43, 44) (Table 1)

Other professions now adopting forms of manual therapy have designated various terms for the subluxation finding. It is this entity that needs to be clinically identified and quantified in consideration with its articular component, in order to resolve the disturbances through specific manipulation. The clinical efficacy of spinal manipulation in addressing the lesion as a cause and effect observation may depend on a range of factors including:

- site
- duration
- severity
- degenerative change(s)
- trauma
- patient age
- hobbies
- pastimes, and
- occupation.

While most VSCs can be corrected or ameliorated, these factors may at times determine grounds for ongoing management or supportive care due to a loss of stability leading to recurrences. Such cases often require specific exercise programs to help strengthen such sites and

minimise exacerbations and relapses, along with lifestyle changes when they are indicated. (3, 4, 7, 12)

**Table 1:** Nominated segmental associations

SEGMENT LEVEL	ASSOCIATED ORGAN/ FUNCTION	AUTHOR	JOURNAL/YEAR
C1/C2 (Rats)	Heart/hypertension	He, Lv, Li, et al (70)	Biomed Res Int. 2017
T2/T3	Raynaud's Disease	Fraser DM (71)	Textbook, 1990
T4	Glove paresthesias	McGuckin N (72,73) DeFranca, Levine	Textbook, 1986 JMPT, 1995
T5	Angina pectoris	Hamberg, Lindall (74)	ACTA Med Scand, 1981
T5 (Rat subjects)	Impaired left ventricular function with enhanced sympathetic support	Lujan Janbaih DiCarlo (75)	J Applied Physiol, 2012
T5/6 (T4-T7)	Gastro-duodenal	Lewit (76)	Textbook, 1985
T6 'Anterior'	Dyspepsia	Rome (77)	Chiropr J Aust, 2000
T6-T9	Duodenal or gastric ulcers	Kameith (78)	Arch Orthop Unfallchir, 1958
[T13 (Rat)	Adrenal	Isa et al (79)	Neuro Sci, 1985
Multiple (Rat)	Depending on level Organ specific	Kimura A, Sato A (80)	Jpn J Vet Res. 1997

As with the diagnosis of headaches, chest pain, short sightedness and psychiatry, a large degree of subjective input may be beneficial to confirm the diagnostic localisation in the case of a VSC. Apart from clinical indications of patients' presenting symptoms and signs, at times patients may also require functional, static or other radiological imaging for further assessment, in order to rule out the possibility of contraindications for physical care. At times, imaging can assist in indicating the most likely area warranting attention as the cause of a patient's symptoms. A thorough case history and examination may comprise a physical palpation, motion palpation, as well as the standard physical, neurological, and orthopaedic testing. The results often correlate with the patients' symptoms as a starting point.

Clinically, a similar rationale with neural implications could be compared to that involved in the reduction of vertebrogenic pain, in cases of intercostal neuralgia, or in subluxation-related sciatica; a condition which can ultimately result in symptoms of paresthesias, dysesthesias, muscle dysfunction of weakness and muscle atrophy. Vertebrogenic conditions cannot logically be

confined solely to musculoskeletal conditions or an 'MSK' model, because of the deep integration and influence with the ANS. (48, 49, 50)

A comorbidity relationship of back pain and certain so-called visceral conditions was noted in 2017 by de Luca and colleagues. They discussed an association of such conditions as diabetes, cardiovascular disorder, obesity, and pulmonary disease with low back pain. The research did not however determine the effect a reduction in the back pain may have had on those conditions. (51)

In more recent times, and contrary to a 1972 opinion proposed by Wyke, Giles discovered free nerve endings in the synovial folds of lumbar vertebral facets. This anatomical contribution tends to focus attention on the disturbed facet being a primary cause of spinal pain, and refocuses more evidence on the importance of restoring normal articular sensory input. (52, 53, 54, 55)

In a detailed presentation, Lee and Salter note chondrocyte mechanotransduction as one of a number of functions at the cellular level which transmits sensory signals to chondrocytes under compressive strain. These histological physical features provide a far more complex picture than just a physical cartilage surface with dysfunction or minimal displacement. (56, 57)

In considering these aspects, it would seem apparent that the subluxation is far more complex than a mere mechanical manipulative lesion. If this was not the case, it would not be independently raised in the literature in association with noted common but apparent neurological symptoms, such as cervicogenic headaches, dyspepsia, blood pressure changes and HRV. See Parts 6, 7, 8 and 10 of this series.

This VSC condition also raises questions regarding the point at which displacement of a conventionally defined subluxated vertebra begins to become clinically significant and at which stage it affects somatosensory and other neuromusculoskeletal symptoms, as well as the mechanism that allows it to develop. The symptoms essentially demonstrate that subluxations involve more complex issues than just physical displacement and dysfunction.

### Vertebrogenic-segment specific

Chiropractic has primarily based its health care model on the neurophysiology of articular segmental spinal influence. (58, 59, 60) Kirpalani and Mitra cited research of the cervical spine and stated that *'each joint produces a distinct referred pain pattern.'* (61,) The efficacy of this is supported by numerous case reports as noted in the literature, most notably the *Index to Chiropractic Literature*. (32)

In 1921, Firth published a chiropractic text with extensive correlations between specific spinal segments and particular clinical findings. Prior to that, in 1910 Palmer had noted a correlation between certain spinal segments and particular visceral and other conditions. (62, 63)

Also one hundred years ago, a medical study by Winsor at the University of Pennsylvania in 1921 conducted autopsies of 75 human and 22 cat subjects. He noted an association of spinal segmental specificity with diseased organs, a 96% correlation rate. In this study, his table of visceral disturbances lists 14 separate conditions. (64)

Warren claims that Winsor set out to disprove the chiropractic hypotheses, only to find his research was consistent with these concepts. (65)

In 1992, Jänig and McLachlan stated that *'The autonomic nervous system supplies each type of target organ via separate pathways which consists of sets or pre-and postganglionic neurones with distinct patterns of reflex activity.'* This has a homeostatic role in the protection and regulation of bodily functions. (66, 67)

It is noted further that neural irritation as explained by Sato and Schmidt, that in relation to the sympathetic reflexes in cats that *'The size of the early reflex component was largest when the*

*afferent volley entered the spinal cord at the same segment or at the segment adjacent to the white ramus.'* (68)

Cramer and Darby note rather specific vertebral correlations with organ nociception through preganglionic autonomic fibres. Their studies support the clinical observation of potential influence through spinal manipulation – the somatovisceral reflex association. They note that *'Since pain is the most important clinical visceral sensation, knowledge of the spinal cord segments to which visceral afferent fibers project (which is the same location as the sympathetic preganglionic cell bodies) is extremely important. This knowledge allows the clinician to more effectively diagnose pathological conditions occurring in the viscera.'* (69)

Stochkendahl and Christensen report *'segmental dysfunction of the cervical and thoracic spine' among other musculoskeletal disorders, as being "under-diagnosed"* as well as a possible cause of thorax pain syndromes. While they recognise that the more serious considerations of chest pain must take priority, some 80% may be benign and approximately 50% of these would be musculoskeletal. The diagnostic and treatment consequences of these facts may at times be complex. (81)

In chiropractic research reported in *Neuroscience Letters*, Desmarais et al demonstrated segmentally responsive somatosympathetic reflexes throughout the thoracic spine. They found that spinal manipulation could modulate the effect of the supraspinal sympathetic reflex skin response to a heat stimulus in healthy humans. (37)

In further research in 1996, Kimura and colleagues confirmed that localised somatic stimulation of spinal-cardiac sympathetic reflexes in rats were segmentally organised.

### Adrenal glands

Isa et al studied anaesthetised rat specimens, where somatic stimulation at the T13 segmental innervation level was activated through pinching, brushing and electric shock to the skin demonstrating a corresponding response of the adrenal glands. They noted *'clear and consistent decreases in blood pressure and renal nerve activity.'* (79, 83)

Sato et al noted an adrenal response with vertebral joint stimulation on rats at spinal levels between T10 and L5. Sato also anaesthetised animals in order to eliminate emotional factors. They found that somatic afferent stimulation can regulate visceral function and that particular organs can at times be segment-specific in regards to gastric motility and urinary vesical contractility. (84, 85, 86)

Others have also noted similar segmental anatomo-physiological associations. Using rat subjects, Strack et al noted that the adrenal gland was supplied with multi-segmental input from sympathetic preganglionic neurons and dorsal root ganglion cells. In acknowledging segmental specificity, they noted that one segment was the dominant source of the innervation. (87)

### Grisel Syndrome

A vertebral subluxation has been recognised as being associated with a relatively rare pathophysiological medical condition of the upper cervical spine called *Grisel Syndrome*. This has been designated as being of uncertain etiology and usually, but not always, presents in children. (88) As a viscerosomatic reflex primary finding, the displacement of a cervical vertebra is adjudged to follow a range of conditions such as acute lymphadenopathy, head-neck surgery, infection, musculoskeletal (cervical ligamentous), physical trauma, and head/neck inflammation. Whether this associated displacement is cause or effect has not been definitively determined, but an apparent neural reflex is noted.

Until relatively recently, the conventional allopathic version of a subluxation would seem to be considered as a purely osseous anatomical displacement with limited consideration of its effects



on associated structures or physiological functions, especially neurological aberrations. Further, this interpretation does not appear to identify or differentiate the degree of displacement which may initiate the onset of symptoms - be they subtle or overt. (89, 90, 91, 92)

While *Grisel Syndrome* is generally considered to be identified by the increase of the C0/C1 pre-dental interspace, Martinez and colleagues identified a case involving a C2/C3 subluxation, while Lopes and Li identified a case at the C3-C4 level. The mechanism of how an infection may initiate a subluxation does not appear to have attracted serious consideration. (93, 94)

This syndrome characterises the integration of particular vertebral subluxations with autonomic and visceral components.

### Scolioses

Lateral spinal curves in scolioses deviate from the midline and range from mild to severe. They may also be subject to VSC dysfunction within those curves. Manipulative management may address those VSCs, but it is also clinically important to restore them towards normal segmental mobility, function for symptomatic amelioration. In addition, maintaining general spinal flexibility within the patient's comfortable limits is also important. (95, 96, 97, 98)

In 1933 and again in 1940, research published by Ussher noted a correlation between particular spinal regions and *simulated condition* of specific organs. This association was related to patterns in lateral spinal curvature. (99, 100)

In a similar vein in 1958, Kameith reported radiographic findings indicating an association between scoliotic orientation and either gastric or duodenal ulcers. He noted that '*the percentile distribution of right-sided and left-sided scolioses coincided with the percentages of duodenal and gastric ulcers...(and further that )...all the scolioses involved vertebral segments corresponding to the stomach and duodenum, that is, T6 to T9.*' Kameith's findings appear to be supported by those of Giampietro et al in 2013, who noted the association of congenital scoliosis and certain health problems. (78, 101)

### Congenital vertebral anomalies

A radiological study by Schey in 1976 examined a relationship between spinal anomalies and visceral abnormalities. While thought to be embryological in origin, there was a high correlation between the particular osseous spinal anomalies and visceral abnormalities. Although not considered vertebrogenic, the vertebro-visceral correlation would be considered relevant in a somato-visceral correlation. (102)

Denton found genitourinary tract and anal anomalies in 53% of cases correlated with lower spine anomalies - particularly sacral. Beals and colleagues noted that 61% of 218 patients with vertebral anomalies exhibited anomalies affecting seven systems. The segmental level only had a mild degree of correlation with the associated diagnosis. (103, 104, 105)

Trenga and colleagues found a 55% correlation of spinal cord anomalies associated with congenital spinal segments. They noted that the more complex the congenital malformation, the greater the incidence of cord anomaly. (106) On the other hand, Theiss and colleagues found minimal correlation between Klippel-Feil syndrome and cervical symptoms. (107)

The identification of block vertebrae by motion palpation is relatively straightforward as noted by Humphreys et al who concluded that '*relatively inexperienced examiners are capable of correctly identifying inter-segmental fixations in the cervical spine.*' However, the subtleties of degree, direction and type of dysfunction and/or minor displacement requires sensitive palpation and appreciation of the minutiae of subluxations. (108)

## Conclusion

The independent and chiropractic evidence presented tends to confirm the model of a pathophysiological role of somato-autonomic factors associated with vertebral subluxations. We could not locate any evidence which contradicted this statement.

Ongoing chiropractic research continues to elucidate how the removal of VSCs produces the necessary benefits, especially at the neuronal level. It would be anticipated that such data would help explain apparent clinical efficacy.



Peter Rome

DC (ret), FICC

[cadaps@bigpond.net.au](mailto:cadaps@bigpond.net.au)

John D Waterhouse

DC, FACC

Private practice, Melbourne

---

**Cite:** Rome P, Waterhouse JD. Neurodynamics of vertebrogenic somatosensory activation and Autonomic Reflexes - a review: Part 5 Diversity in Vertebral Subluxations. Asia-Pacific Chiropr J. 2021;1.4. URL [apcj.net/papers-issue-2-4/#RomeWaterhousePart5Diversity](https://apcj.net/papers-issue-2-4/#RomeWaterhousePart5Diversity)

## References

1. Branyon B. Healing hands: using osteopathic manipulative treatment to address visceral structures through somatovisceral reflexes: a case study in gastroesophageal reflux disease. J Am Acad Osteop. 2008;18(4):29-31.
2. DeStefano L. Greenman's principles of manual medicine. 5th ed. Philadelphia. Lippencott Williams and Wilkins. 2016.
3. Gatterman MI. Foundations of chiropractic subluxation. 2nd edn. St Louis: Elsevier Mosby;2005
4. Haldeman S. Principles and practice of chiropractic. 3rd edn. New York: McGraw-Hill Medical: 2004.
5. King HH, Jänig W, Patterson MM. Segmental and suprasegmental mediation of somatosensory interactions. In: The science and clinical application of manual therapy. Edinburgh: Churchill Livingstone.2011:108-191.
6. King HH. Impact of manual therapy on systemic disorders and physiologic functions. In: Research related to clinical applications of manual therapy for musculoskeletal and systemic disorders from osteopathic experience. (p 208) In: King196-216. <https://musculoskeletalkey.com/research-related-to-clinical-applications-of-manual-therapy-for-musculoskeletal-and-systemic-disorders-from-the-osteopathic-experience/>
7. Leach RA The chiropractic theories: principles and clinical application. 3rd edn. Baltimore: Williams & Wilkins 1994
8. Painter FM. What is the chiropractic subluxation? <https://chiro.org/Subluxation/>
9. Rosner AL. The role of subluxation in chiropractic. Des Moines. Foundation for Chiropractic Education and Research. 1997
10. Slosberg M. Validating chiropractic. Cutting-edge research to improve patient outcomes. <https://cdn.vortala.com/childsites/uploads/41/files/validating-chiropractic.pdf> (Extracts)
11. Slosberg M. Applying cutting-edge research to everyday practice. Chiropractic and exercise to improve physical well-being and brain function. 2014. 91pps.
12. Henderson CN. The basis for spinal manipulation: chiropractic perspective of indications and theories. J Electromyogr Kinesiol. 2012;22(5):632-642.
13. Cramer GD, Henderson CNR, Little JW, Daley C, Grieve TJ. Zygapophyseal joint adhesions after induced hypomobility. J Manipulative Physiol Ther. 2010;33(7):508-518.



14. Dishman RW. Static and dynamic components of the chiropractic subluxation complex: a literature review. *J Manipulative Physiol Ther.* 1988;11(2):98-107.
15. Seaman DR, Winterstein JF. Dysafferentation: a novel term to describe the neuropathophysiological effects of joint complex dysfunction. A look at likely mechanisms of symptom generation. *J Manipulative Physiol Ther.* 1998;21(4):267-80.
16. Leach RA. (7, p 89-119)
17. Beck R. Functional neurology for practitioners of manual medicine. London. Elsevier Health Sciences. 2011:321.
18. Mootz RD. Theoretical models of subluxation. Chapter 10. In: Gatterman MI. Foundations of chiropractic subluxation. 2nd edn. St Louis: Elsevier Mosby;2005;227-244. Also at: Mootz RD. Theoretical models of subluxation. Chapter 10. In: Physical medicine and rehabilitation. 2016. <https://musculoskeletalkey.com/theoretic-models-of-subluxation/>
19. Kent C. Models of vertebral subluxation: a review. *J Vert Subluxation Res.* 1996;1(1):1-7
20. Gongal'skiĭ, VV, Moroz NF. [Vertebral joint meniscoids in the norm and in functional blocking of the vertebral segment. *Lik Sprava.* 2002;2:96-99.
21. Leach RA. (7) p 97
22. Tozzi P. A unifying neurofasciogenic model of somatic dysfunction – underlying mechanisms and treatment. *J Bodyw Mov Ther.* 2015;19(2):310-26.
23. Heilig D, The thrust technique. *J Am Osteop Assoc.* 1981;81(4):244-8
24. Garcia F, Forrester B. The neuro sensorimotor integrator. Carrick Institute. <https://carrickinstitute.com/citv-technology-review-the-nsi-unit/>
25. Sato A, Sato Y, Schmidt RF. The impact of somatosensory input on autonomic functions. In: Blaustein MP, Grunicke H, Pette D, Schultz G, Schweiger M. (eds). *Rev Physiol Biochem Pharmacol.* Berlin: Springer;1997;130:1-2,138,231,258
26. Lantz CA. Immobilization degeneration and the fixation hypothesis of chiropractic subluxation. *Chiropr Research J* 1988;1(1):21-46
27. Leach RA. (7) p 214-5
28. Fryer G, Morris T, Gibbons P. Paraspinal muscles and intervertebral dysfunction: part two. *J Manipulative Physiological Ther.* 2004;27:348-57.
29. Senzon S. The chiropractic vertebral subluxation. Parts 1 -10. *J Chiropr Humanit.* 2018;25C:10-168,
30. Rome PL. Medical evidence recognising the vertebral subluxation complex. *Chiropr J Aust.* 2016;44(4):304-307
31. Rome PL, Waterhouse JD. Evidence informed vertebral subluxation - a diagnostic and clinical imperative. *J Philos Principles Prac Chiropr.* 2019; Dec 2;12-34
32. Index to Chiropractic Literature. [https://www.chiroindex.org/?advanced\\_search=1#results](https://www.chiroindex.org/?advanced_search=1#results)
33. Osteopathic Medical Digital Repository. <https://ostemed-dr.contentdm.oclc.org/digital/search/advanced>
34. Ernst E. Subluxation, a myth that is deeply ingrained in the chiropractic mind-set. Feb 2015. <https://edzardernst.com/2015/02/subluxation-a-myth-that-is-deeply-ingrained-in-the-chiropractic-mind-set/>
35. WHO guidelines on basic training and safety in chiropractic. 2005 <https://www.who.int/medicines/areas/traditional/Chiro-Guidelines.pdf>
36. Sato A, Sato Y, Schmidt RF.(25) p 138
37. Desmarais A, Descarreaux M, Houle S, Piché M. Tuning the gain of somato-sympathetic reflexes by stimulation of the thoracic spine in humans. *Neurosci Lett.* 2011;490(2):107-111. doi: 10.1016/j.neulet.12.037
38. Keller TS, Colloca CJ, Moore RJ. Increased multi-axial lumbar motion responses during multi-impulse mechanical force manually assisted spinal manipulation. *Chiropr Osteop* 2006;14:6. doi:10.1186/1746-1340-14-6
39. Colloca CJ, Keller TS, Harrison DE, et al. Spinal manipulation force and duration affect vertebral movement and neuromuscular response. *Clin Biomech.* 2006;21(3):254-62
40. Burns L, Chandler LC, Rice RW. Pathogenesis of visceral disease following vertebral lesions. *J Am Osteop Assoc, Chicago* 1948. (Note Dr Burns has also published at length on her extensive research, particularly in the *J Am Osteop Assoc* and the *AT Still Research Institute Bulletin*.)
41. Cleveland CS. Researching the subluxation of the domestic rabbit: a pilot research program conducted at the Cleveland Chiropractic College. *Pub Cleveland Chiropractic College, Kansas City, Missouri*, 24pps. (See also *Sci Review Chiropr.* Aug 1965;(4):5-28 .
42. Rosner A. Getting down to brass tacks: The neurophysiology of spinal manipulation. *Dynamic Chiropr.* 2015;8(9): [http://www.dynamicchiropractic.ca/mpacms/dc\\_ca/columnist\\_other\\_articles.php?id=1591](http://www.dynamicchiropractic.ca/mpacms/dc_ca/columnist_other_articles.php?id=1591).
43. Cramer G, Henderson C, Khalsa P, Pickar J. Animal models. In: Basic science research related to chiropractic spinal adjusting: the state of the art and recommendations. *J Manipulative Physiol Ther* 2006;29(9):726-761.
44. Henderson CNR. Animal models in the study of subluxation and manipulation: 1964-2004. In: Gatterman MI. (3) p-47-103.
45. Rome PL. Terminology relating to the vertebral subluxation complex and the manipulative sciences. (Part 1) *Chiropr J Aust* 2017;45(2):73-89

46. Rome PL. Terminology relating to the vertebral subluxation complex and the manipulative sciences. (Part 2) *Chiropr J Aust* 2017;45(2):90-130
47. Rome PL. Usage of chiropractic terminology in the literature: 296 ways to say "subluxation". Complex issues of the vertebral subluxation. *Chiro Tech* 1996;8(2):49-60
48. Pickar JG. Neurophysiological effects of spinal manipulation. *Spine J* 2002;2(5):357-371,
49. Colloca CJ, Keller TS, Gunzburg R, Vandeputte K, Fuhr AW. Neurophysiologic response to intraoperative lumbosacral spinal manipulation. *J Manipulative Physiol Ther.* 2000;23(7):447-457
50. Colloca CJ, Keller TS, Gunzburg R, Neurophysiologic characterisation of in vivo lumbar spinal manipulation Part II. Neurophysiological response. *J Manipulative Physiol Ther.* 2003;26(9):579-547
51. de Luca KE, Parkinson L, Haldeman S, Byles JE, Blyth F. The relationship between spinal pain and comorbidity: a cross-sectional analysis of 579 community-dwelling, older Australian women. *J Manipulative Physiol Ther* 2017;40(7):459-466
52. Wyke B. Articular neurology – a review. *Physiotherapy* 1972;58(3):94-99
53. Giles LGF. Back off, memoir of the vicissitudes of a complementary health practitioner. San Francisco. *Blurb.com*.2017:14-20
54. Giles LG, Tayloe JR, Cockson A. Human zygapophyseal joint synovial folds. *ACTA Anat (Basel).* 1986;126(2):110-114.
55. Giles LG, Harvey AR. Immunohistochemical demonstration of nociceptors in the capsule and synovial folds of human zygapophyseal joints. *Br J Rheumatol* 1987;26(5):362-365
56. Lee H-S, Salter DM. Biomechanics of cartilage and osteoarthritis. 2015. <http://dx.doi.org/10.5772/60011> <https://cdn.intechopen.com/pdfs-wm/48287.pdf> P 45
57. Zimny ML. Mechanoreceptors in articular tissues. *Am J Anat* 1988;182(1):16-32. <http://www.biology-pages.info/M/Mechanoreceptors.html#Proprioception>.
58. Dwyer A, Aprill C, Bogduk N. Cervical zygapophyseal joint pain patterns. I: A study of normal volunteers. *Spine.* 1990;15(6):453-7.
59. Aprill C, Dwyer A, Bogduk N. Cervical zygapophyseal joint pain patterns. II: A clinical evaluation. *Spine.* 1990;15(6):358-61.
60. Dreyfuss P, Tibiletti C, Dwyer CJ. Thoracic zygapophyseal joint pain patterns. A study in normal volunteers. *Spine.* 1994;19(7):807-811.
61. Kirpalani D, Mitra R. Cervical facet joint dysfunction: a review. *Arch Phys Med Rehab.* 2008;89:770-4
62. Firth JN. A textbook on chiropractic symptomatology 2nd edn. Self Published 1921.
63. Palmer DD. The science, art and philosophy of chiropractic. Portland: Portland Printing House Co. 1910:913-970
64. Winsor H. Sympathetic segmental disturbances. The evidence of the association in dissected cadavers of visceral disease with vertebrae deformities of the same sympathetic segment. *New York Med Times* Nov 1921;XLIX(ii):267-271. [Also summary <http://www.dorn-method.com/pdf/WinsorMDArticleSympatheticSegmentalDisturbances.pdf>]
65. Warren J. Could 50 autopsies in 1921 have revealed the keys to long life? <http://drjaywarren.com/windsors-autopsies/> . 2017
66. Jänig W, McLachlan EM. Specialized functional pathways are the building blocks of the autonomic nervous system. *J Autonom Nerv Syst* 1992;41(1-2):3-13
67. Jänig W, Häbler HJ. Specificity in the organization of the autonomic nervous system: a basis for precise neural regulation of homeostatic and protective body functions. *Prog Brain Res* 2000;122:351-367
68. Sato A, Sato Y, Schmidt RF. (25) p78.
69. Cramer GD, Darby S. Basic and clinical anatomy of the spine, spinal cord, and ANS. St Louis: Mosby;1995: Table 1. p342 .
70. He Z-B, Lv Y-K, Li H, et al. Atlantoaxial misalignment causes high blood pressure in rats: a novel hypertension model. *Biomed Res Int* 2017;16: 5986957
71. Fraser DM. T3 Syndrome. In: Paterson JK, Burns L, editors. *Back Pain - an international review*. Dordrecht: Kluwer Academic, 1990;387-390
72. McGuckin N. The T4 Syndrome. In: Grieve GP, ed. *Modern manual therapy of the vertebral column*. New York: Churchill Livingstone, 1986;370-376
73. DeFranca GG, Levine LJ. The T4 Syndrome. *J Manipulative Physiol Ther* 1995;18(1):34-37.
74. Hamberg J, Lindall O. Angina pectoris symptoms caused by thoracic spine disorders. *Acta Med Scand Suppl* 1981;644:84-86
75. Lujan HL, Janbair H, DiCarlo SE. Dynamic interaction between the heart and its sympathetic innervation following T5 spinal cord transection. *J Applied Physiol* 2012;113(8):1332-1341.
76. Lewit K. Manipulative therapy in rehabilitation of the locomotor system. London: Butterworths, 1985;339-340.
77. Rome PL. Anterior T6 subluxation syndrome: neurospinal dysfunction within a vertebral subluxation complex. *Chiropr J Aust.* 2000;30(4):127-137.
78. Kameith H. [The thoracic spine as a pathogenic factor.] *Arch Orthop Unfallchir.* 1958;49(6):585-606. (German) (Pubmed English extract) (Cited as 'Pathogenic importance of the thoracic portion of the vertebral column.' *JAMA.* 1958;Nov 15:1586.)

79. Isa T, Kurosawa M, Sato A, Swenson RS. Reflex responses in the adrenal sympathetic nerve to electrical stimulation of somatic afferent nerves in the rat. *Neurosci Res.* 1985;3(2):130-44.
80. Kimura A, Sato A. Somatic regulation of autonomic functions in anesthetised animals - neural mechanisms of physical therapy and acupuncture. *Jpn J Vet Res.* 1997;45(3):137-145.
81. Stochkendahl MJ, Christensen HW. Chest pain in focal musculoskeletal disorders. *Med Clin North Am.* 2010;94(2):259-273. doi: 10.1016/j.mcna.2010.01.007.
82. Kimura A, Sato A, Sato Y, Suzuki H. A- and C-reflexes elicited in cardiac sympathetic nerves by single shock to a somatic afferent nerve include spinal and supraspinal components in anesthetised rats. *Neurosci Res.* 1996;25(1):91-96.
83. Sato A, Sato Y, Schmidt RF. (25) p 219-253.
84. Sato A. Neural mechanisms of somatic sensory regulation of catecholamine secretion from the adrenal gland. *Adv Biophys* 1987;23:39-80.
85. Sato, A., Sato, Y., Schmidt, RF : Catecholamine secretion and adrenal nerve activity in response to normal and inflamed knee joints in cats. *J. Physiol* 1986;375: 611-624
86. Sato A, Sato Y, Schmidt RF. The effects of knee joint stimulation on the sympatho-adrenal medullary functions in anesthetized cats. *Neurosci Res* 1985;3(Suppl 1):S18.
87. Strack AM, Sawyer WB, Marubio LM, Loewy AD. Spinal origin of sympathetic preganglionic neurons in the rat. *Brain Res* 1988;455(1):187-191
88. Park SH, Park SH, Lee SH. Grisel syndrome: pathophysiological evidence from magnetic resonance imaging findings. *Ann Rehabil Med.* 2013;37(5):713-716.
89. Dewan K, Giannoni C. Radiology quix case 2. Nontraumatic atlantoaxial subluxation (also known as Grisel syndrome. *Arch Otolaryngol Head Neck.* 2012;1238(2):199-201.
90. Harma A, Firat Y. Grisel syndrome: nontraumatic atlantoaxial rotatory subluxation. *J Craniofac Surg* 2008;19(4):119-1121
91. Ortega-Evangelio G, Alcon JJ, Alvarez-Pitti J, Juncos M, Lurbe E. Eponym: Grisel syndrome. *Eur J Pediatr.* 2011;170(8):965-968.
92. Yochum TR, Rowe LJ. Essentials of skeletal radiology. Vol 1. 2nd ed. Baltimore. Williams & Wilkins. 1996:148
93. Martínez-Lage JF1, Morales T, Fernandez Cornejo V. Inflammatory C2-3 subluxation: a Grisel's syndrome variant. *Arch Dis Child.* 2003 Jul;88(7):628-9.
94. Lopes DK, Li V. Midcervical postinfectious ligamentous instability: a variant of Grisel's syndrome. *Pediatr Neurosurg* 1998;29:133-137
95. Morningstar MW, Stitzel CJ, Siddiqui A, Dovorany B. Chiropractic treatments for idiopathic scoliosis: a narrative review based on SOSORT outcome criteria. *J Chiropr Med* 2017;16(1):64-71.
96. Byun S, Han D. The effect of chiropractic techniques on the Cobb angle in idiopathic scoliosis arising in adolescence. *J Phys Ther Sci* 2016;28(1):1106-1110.
97. Poussa M, Mellin G. Spinal mobility and posture in adolescent idiopathic scoliosis at three stages of curve magnitude. *Spine* 1992;17(7):757-760
98. Senna MK, Machaly SA. Does maintained spinal manipulation therapy for chronic nonspecific low back pain result in better long-term outcome? *Spine* 2011;36(18):1427-1437.
99. Ussher NT. Spinal curvatures - visceral disturbances in relation thereto. *Calif Western Med* June 1933,38(16):423-428.
100. Ussher NT. The viscerospinal syndrome - a new concept of visceromotor and sensory changes in relation to deranged spinal structures. *Annals Internal Med.* 1940;13(2):2057-2090.
101. Giampietro PF, Raggio CL, Blank RD, et al. Clinical, genetic and environmental factors associated with congenital vertebral malformations. *Mol Syndromol* 2013;4(1-2):94-105
102. Schey WL. Vertebral malformations and associated somatovisceral abnormalities. *Clin Radiol* 1976;27:341-353.
103. Denton JR. The association of congenital spinal anomalies with imperforate anus. *Clin Orthop Relat Res* 1982;162:91-98.
104. Beals RK, Robbins JR, Rolfe B. Anomalies associated with vertebral malformations. *Spine* 1993;18(10):1329-1332
105. Rai AS, Taylor TKF, Smith GHH, Cumming RG, Plunkett-Cole M. Congenital abnormalities of the urogenital tract in association with congenital vertebral malformations. *Bone Joint J* 2002;84(6):891-895.
106. Trenga AP, Sungla A, Feger MA, Abel MF. Patterns of congenital bony spinal deformity and associated neural anomalies on x-ray and magnetic imaging. *J Child Orthop* 2016;10(4):343-362
107. Theiss SM, Smith MD, Winter RB. The long-term follow-up of patients with Klippel-Feil syndrome and congenital scoliosis. *Spine* 1997;22(11):1219-1222.
108. Humphreys BK, Delahaye M, Peterson CK. An investigation into the validity of cervical spine motion palpation using subjects with congenital block vertebrae as a 'gold standard'. *BMC Musculoskelet Disord.* 2004 Jun 15;5:19. doi: 10.1186/1471-2474-5-19.